

Insecticide Resistance Management Guidelines for Beet Armyworm in Lettuce

John C. Palumbo


University of Arizona, Department of Entomology

in collaboration with the **Insecticide Resistance Action Committee (IRAC)**, a CropLife specialist technical group

The figures below illustrate insecticide options available for chemical management of beet armyworm and other important lepidopterous larvae during the growing season. **Figure 1** provides a relative index of efficacy for insecticides currently labeled on lettuce for management of beet armyworm. The index is based on empirical data generated from local field trials. **Figure 2** offers guidance for each insecticide product and its most effective fit at various crop stages throughout the crop season.

These charts should serve as a guide to PCAs and growers for avoiding the overuse of a single product based on its IRAC defined mode of action (MOA), and as a reference for selecting products/MOAs with which to rotate throughout the season for the purpose of maximizing and sustaining product efficacy. This management approach should not be difficult to implement given the number of insecticide products with distinctly different MOA available for management of lepidopterous larvae throughout the season (Fig 1 and 2).

Figure 1.
Lepidopterous Larvae Management in Desert Lettuce Crops



Relative Efficacy Index For Lep Larvae in Desert					
Product	IRAC ¹ MOA	Beet armyworm	Cabbage loopers	Corn earworm	Comments*
Lannate	1A	**	*	***	Tank mix with another product for broad spectrum Lep activity; provides thrips control; PHI: 10 d on lettuce; 7 d spinach
Acephate	1B	*	**	**	Tank mix with another product for broad spectrum Lep activity; PHI: 14-21 d on head lettuce, 7 d on cauliflower ; provides thrips control
Endosulfan	2A	*	***	**	Tank mix with another product for broad spectrum Lep activity; PHI: 21 d for head lettuce and celery ; 7 d on cauliflower
Pyrethroids	3	*	***	***	Tank mix with another product for broad spectrum Lep activity; use high labeled rates; PHI: varies with products
Radiant	5	***	***	***	Stand alone Lep, leafminer, and thrips control; PHI: 1 day on leafy veg and Brassica crop groups
Proclaim	6	***	**	***	Stand alone Lep control; a penetrating adjuvant may enhance residual control; PHI: 7 day on leafy vegetable and Brassica head and stem crop groups
Bt (i.e. Dipel)	11B	*	**	*	Tank mix with another product for broad spectrum Lep activity; numerous Bt products available; PHI: 0 d -good spray coverage desirable
Intrepid	18A	***	***	**	Tank mix with another product for broad spectrum Lep activity; PHI: 1 day on leafy vegetable and Brassica crop groups -good spray coverage desirable
Avaunt	22	***	***	**	Tank mix with another product for broad spectrum Lep activity; PHI: 1 day on leafy vegetable and Brassica crop groups -good spray coverage desirable
Synapse	28	***	***	***	Stand alone Lep control; PHI: 1 day on leafy vegetable and Brassica leafy crop groups
Coragen	28	***	***	***	Stand alone Lep and leafminer control; PHI: 1 day for Leafy Veg crop group; 3 d for Brassica leafy crop group for both soil and foliar uses
Voliam Xpress	28+3	***	***	***	Stand alone Lep and leafminer control; PHI: 1 day for head and leaf lettuce; 3 d for Brassica head and stem crop group.
Volium Flexi	28+4A	***	***	***	Stand alone Lep and leafminer control; PHI: 7 day for leaf veg crop group; 3 d for Brassica head and stem crop group. Has aphid activity.
Durivo	28+4A	***	***	***	Stand alone Lep and leafminer control; PHI: 30 day for leaf veg and Brassica crop groups; Has aphid activity.
Vetiva	28+16	***	***	***	Stand alone Lep control; PHI: 7 day for Leafy Veg crop group; 1 d for Brassica leafy crop group. Has whitefly activity.
***	Good residual control (7-14 d)				
**	Marginal residual control (4-6 d)				
*	Poor residual control (1-3 d)				

¹ IRAC Mode of Action - for more info go to - <http://www.irac-online.org/>
 * always consult the label before applying any of these products

Figure 2.

UA IPM Guidelines for Lep Management in Leafy Vegetables

Alternatives for Lep Larvae Control by Crop Stages

Insecticide	IRAC MOA	Soil – at plant	Stand establishment		Thinning to Heading			Heading to Harvest		
			Coty-1 leaf	2-4 leaf	5-8 leaf	9-15 leaf	15-20 leaf	Pre - head	Early heading	2-4" head
Radiant	5		Green	Green	Green	Green	Green	Green	Green	Green
Proclaim	6		Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow	Yellow
Intrepid	18				Blue	Blue	Blue	Blue	Blue	
Avaunt	22A				Light Blue	Light Blue	Light Blue	Light Blue	Light Blue	
Coragen	28	Red	Red	Red	Red	Red	Red	Red	Red	Red
Durivo	28+4A	Red	Red	Red						
Voliam Xpress	28+3		Red	Red	Red	Red	Red	Red	Red	Red
Voliam Flexi	28+4A			Red	Red	Red	Red	Red	Red	
Synapse	28			Red	Red	Red	Red	Red	Red	
Vetica	28+16			Red	Red	Red	Red	Red	Red	
Lannate	1A		Light Red	Light Red	Light Red	Light Red	Light Red	Light Red	Light Red	Light Red
Orthene	1B			Light Red	Light Red	Light Red				
Endosulfan	2A			Light Blue	Light Blue	Light Blue				
Pyrethroids	3		Grey	Grey	Grey	Grey	Grey	Grey	Grey	Grey
Bt	11B				Dark Green	Dark Green	Dark Green			

*** Minimum of 4 effective MOA Effectives at any crop stage

Additional tactics should be practiced to avoid the development of resistance by beet armyworm to any of these products/MOA as follows:

- Apply insecticides only when needed. Time insecticide applications based on UA recommended action thresholds (<http://ag.arizona.edu/crop/>).
- Ideally, the management strategy that presents the lowest risk to insecticide resistance is one where consecutive applications of the same product/MOA **are not** made in the same lettuce field.
- This can be achieved by rotating to an alternative product/MOA on each subsequent spray application to eliminate consecutive uses of the same MOA (see examples in **Figure 3-5** below). Whenever possible, consider using any single product/MOA only once per lettuce field per crop season.
- In lettuce fields where a product/MOA is required more than once, limit the total usage of that product/MOA to 2 applications per field per crop season. (i.e., no more than 2 uses of any IRAC MOA or insecticide with the same color code), and avoid using it on consecutive applications.
- Use only recommended products and rates necessary to accomplish desired control (Fig 1 and 2).
- Do not apply any active ingredient below labeled rates as this may result in poor product performance, unacceptable insect damage and an increased risk of resistance.
- Apply insecticides by directed ground sprays to optimize spray deposition and coverage whenever possible.
- Do not apply tank-mixtures containing 2 or more of the newer chemistries (IRAC Groups - 5, 6, 18, 22 and 28) when controlling lepidopterous larvae. Not only is this expensive, but generally not necessary based on past performance trials (Fig 1).

Specific resistance management recommendations have been developed for the Diamides (IRAC group 28) for *beet armyworm* on lettuce crops grown in the western U.S. Given the residual effectiveness of these compounds, along with their flexibility in application, it will be important to adhere to the guidelines below when using Diamide products as an effort to sustain the efficacy of this new class of insecticide chemistry.

- The Diamide products (IRAC Group 28) offer flexibility in application; they can be applied to plant foliage translaminarily through foliar sprays, or systemically via soil applications.
- If a Diamide product is applied as a foliar spray, consider using this MOA only once per lettuce field per crop season. If a Diamide spray is required more than once, limit the total usage to 2 foliar spray per field and do not use them in consecutive applications (Figure 3).
- **Do not** apply a foliar Diamide spray **prior to** or **following** the use of a soil application of chlorantraniliprole (Figure 4 and 5).
- If a Diamide product is soil applied prior-to or at-planting, as an in-furrow spray or shank injection, **do not spray** a Diamide product on that crop at any time during the remainder of the crop season (Figure 4).
- If a Diamide product (IRAC Group 28) is applied as a post-emergence treatment through drip irrigation, **do not spray** any Diamide products on that crop prior to the Diamide chemigation, or at any time thereafter during the crop season. (Figure 5).
- Do not apply more than **1** application of a Diamide product to the soil regardless if chemigated through drip irrigation or soil applied at planting. If additional beet armyworm control is needed during the crop season, use a non-Diamide foliar alternative. See Figures 1 and 2 for alternative products/MOA.
- Consider using an adjuvant with foliar Diamide applications to assist in spray atomization and penetration, and to provide uniform deposition of spray droplets on foliage; this is particularly important in cole crops.
- In areas where alfalfa is grown in proximity to lettuce, **do not** apply any Diamide product (Coragen, Voliam Xpress) in alfalfa at any time.
- In areas where cotton is grown in proximity to lettuce, **do not** apply any Diamide product (Coragen) in cotton at any time.
- **Do not use** any soil or foliar applied Diamide product on nursery grown plants (e.g., cabbage or cauliflower) destined for field transplanting.

Figure 3




Figure 4




Figure 5

