## **Lepidopterous Larvae Management in Desert Produce Crops, 2024**

Poor residual control



	IRAC 1	Beet	Cabbage	Corn	Diamondback	
Product	MOA	armyworm	looper	earworm	moth **	Comments*
Lannate	1A					Tank mix with another product for broad spectrum Lep activity; provides
Lailliate	17					thrips control; PHI: 10 d on lettuce; Has activity against DBM adults
Acephate	1B					Tank mix with another product for broad spectrum Lep activity; PHI: 21 d or
						head lettuce, celery; 14 d brussels sprouts and cauliflower.
Dibrom	1B					Short residual product. For DBM use 2 pints / acre. Has good adult DBM
	<u> </u>					activity also. PHI: 7-d on labeled cole crops.
Pyrethroids	3					Tank mix with another product for broad spectrum Lep activity; PHI: varies
	<b> </b>					with products; Has shown adult and larval DBM activitiy. Stand alone Lep, leafminer, and thrips control; PHI: 1 day oncole crops and
Radiant/Entrust	5					leafy byeges; Use rates at 5-7 oz depending on pest spectrum.
	1					Stand alone Lep control; PHI: 7 day on cole and leafy veg crops; use at rates
Proclaim	6					above 3.6 oz; if cabbage looper present tank-mixe with a pyrethroid.
	1					Tank mix with another product for broad spectrum Lep activity, numerous B
Bt (Dipel/Xentari)	11B					products available; PHI: 0 d; use rates of 1.5 lb or > for DBM
_	1					Tank mix with a pyrethroid for cabbage looper. Apply at 12 oz for best
Cormoran	15+4A					residual. Has whitefly adult knockdown activity. PHI: 7-d on cloe crops.
Intropid	101					Tank mix with another product for broad spectrum Lep activity; PHI: 1 day;
Intrepid	18A					good spray coverage desirable; use 16 oz rate for DBM
Avaunt	22					Tank mix with another product for broad spectrum Lep activity; PHI: 1 day,
Avaulit	22					use higher rates for best control of all Leps.
Coragen	28					Stand alone Lep and leafminer control; PHI: 1 day for Leafy veg; 3-d for cole
						crops. Avoid use for DBM; resistance has been detected.
Exirel	28					Foliar only; Stand alone Lep, whitefly and leafminer control; PHI: 1 day for
	<u> </u>					Leafy veg annd cole crops - Use at or above 15 oz for DBM.
Verimark	28					Soil only; Stand alone Lep, whitefly and leafminer control; Use at 13.5 oz.
	1					for best residual effectiveness. PHI: N/A Stand alone Lep and leafminer control; PHI: 1 day for Leafy veg annd cole
Harvanta	28					crops - Use at 16.9 oz rate for DBM.
	1					Stand alone Lep and leafminer control; PHI: 1 day forlettuce; 3 day for cole
Besiege	28+3					crops. Avoid use for DBM; resistance has been detected.
						Soil only; Stand alone Lep and leafminer control; PHI: 30 day for leafy and
Durivo	28+4A					cole crops; Avoid use for DBM; resistance has been detected.
Minasta Dua	20.6					Stand alone Lep, whitefly and leafminer control; PHI: 7 day for leafy
Minecto Pro	28+6					vegetables/celery. Use 10 oz rate. Not labeled on cole crops or spinach.
	Good resi	idual control (7-14	d); For DBM - 7 d	residual	1 IRAC Mode of A	ction - for more infor go to - http://www.irac-online.org/
	Suppression					the label before applying any of these products

\*\* Based on 2019-2023 research on local DBM populations