

Whitefly Management in Fall Melons - 2024



Relative Efficacy Index

Product	IRAC ¹ MOA	Adults	Nymphs	
Vydate, Lannate	1A	••	•	Provides knockdown control (1-3 days) of adults when tank-mixed with a pyrethroid Bifenthrin or Danitol
Imidacloprid	4A	•	•••	Generics available; Soil: use highest rate for residual nymph control (0.375 lb ai/ac); may not provide adequate adult control; does not provide consistent performance as post-emergence side dress.
Platinum	4A	•	•••	Soil: use highest rate for residual whitefly control; can be effective as post-plant, side dress or chemigation through drip, marginal residual control when applied at-planting in produce crops due to high water solubility
Assail	4A	•••	•••	Tank-mix with a pyrethroid for improved adult knockdown. Use at high rates (8 oz of Assail 30SG); Relatively safe against honeybees
Venom, Scorpion	4A	•••	•••	<u>Foliar</u> : Foliar applications provide excellent adult knockdown at high rates. <u>Soil</u> : effective as at-planting or post-plant side dress/chemigation treatment on fall melons. Use at high rates.
Transform	4C	•••	••	Provides excellent knockdown (1-3 days) and fair residual control (7 d) of adults ; Provides good suppression of CYSDV; Will also provide good control of Lygus. PHI: 1 day
Sivanto	4D	•••	•••	For <u>soil</u> use only on cantaloupes/honeydews; apply at 28 oa at-plant. 21 D PHI; <u>Foliar and soil</u> uses on watermelons. Apply at higher label rate (10-14 oz) as foliar spray; can provide good knockdown of adults
Durivo	4A+28	••	••	Soil : Contains chlorantraniliprole; will control loopers and beet armyworms; Use at higher rates; effective as at-planting but has shown to provide marginal residual control in produce crops due to high water solubility
Cormoran	4A+15	••	••	Contains acetamiprid+novaluron. Has fair-good residual activity against nymphs and adults. PHI: 7 days.
Knack	7	•	•••	Apply at higher rates for residual control; slow acting. Does not provide adult knockdown control; Will eventually suppress adults numbers by cycling out the immature population.
PQZ	9B	•••	•••	Provides excellent knockdown (1-3 days) and residual control of adults and nymphs; Provides good suppression of CYSDV; Relatively safe against honeybees; PHI: 1 day
Sefina	9D	•••	•••	Provides good knockdown (1-3 days) and residual control of adults and nymphs; Provides good suppression of CYSDV; Relatively safe against honeybees; PHI: 0 day
Courier	16	•	•••	Apply at higher rates for residual control (13.6 oz/ac); slow acting. Does not provide adult knockdown control; Will eventually suppress adults numbers by cycling out the immature population.
Oberon	23	•	•••	Apply at higher rates for residual control (8.5 oz); slow acting. Does not provide adult knockdown control; Will eventually suppress adults numbers by cycling out the immature population.
Exirel	28	•••	•••	Apply at higher rates for longer residual control (15-20 oz); a little slow; may take up to 3 days to observed adult knockdown, but should provide good residual control of nymphs and adults. Use a penetrating adjuvant.
Minecto Pro	28+6	•••	•••	Apply at high rate (10 oz); a little slow; may take up to 3 days to observed adult knockdown, but should provide good residual control of nymphs/adults. Use a penetrating adjuvant. Active against spider mites.

•••	Good residual control
••	Fair-Marginal residual control
•	Poor control

¹ IRAC Mode of Action - for more info go to - <http://www.irac-online.org/>

* always consult the label before applying any of these products, particularly when pollinators are present.